

7.8 Best approximation and least squares

We wish to find the best approximation to the solutions.

Minimum distance problem

The Minimum Distance Problem in R^n Given a subspace W and a vector \mathbf{b} in R^n , find a vector $\hat{\mathbf{w}}$ in W that is closest to \mathbf{b} in the sense that $\|\mathbf{b} - \hat{\mathbf{w}}\| < \|\mathbf{b} - \mathbf{w}\|$ for every vector \mathbf{w} in W that is distinct from $\hat{\mathbf{w}}$. Such a vector $\hat{\mathbf{w}}$, if it exists, is called a *best approximation to \mathbf{b} from W* (Figure 7.8.1).

- Answer:

Theorem 7.8.1 (Best Approximation Theorem) If W is a subspace of R^n , and \mathbf{b} is a point in R^n , then there is a unique best approximation to \mathbf{b} from W , namely $\hat{\mathbf{w}} = \text{proj}_W \mathbf{b}$.

- Distance from \mathbf{b} to a subspace W .
- $d = \|\mathbf{b} - \text{proj}_W(\mathbf{b})\| = \|\text{proj}_{W^c}(\mathbf{b})\|$.

Least square solutions of the linear system.

- $A\mathbf{x}=\mathbf{b}$.
- If this is inconsistent, minimize $\|\mathbf{b}-A\mathbf{x}\|$.

Definition 7.8.2 If A is an $m \times n$ matrix and \mathbf{b} is a vector in R^m , then a vector $\hat{\mathbf{x}}$ in R^n is called a *best approximate solution* or a *least squares solution* of $A\mathbf{x} = \mathbf{b}$ if

$$\|\mathbf{b} - A\hat{\mathbf{x}}\| \leq \|\mathbf{b} - A\mathbf{x}\| \quad (5)$$

for all \mathbf{x} in R^n . The vector $\mathbf{b} - A\hat{\mathbf{x}}$ is called the *least squares error vector*, and the scalar $\|\mathbf{b} - A\hat{\mathbf{x}}\|$ is called the *least squares error*.

Finding the least squares solutions of linear systems.

- $Ax=b$.
- Ax is in $\text{col}(A)$.
- $\|b-Ax\|$ is minimized when $Ax=\text{proj}_{\text{col}(A)}b$.
- This is consistent. Every system has the least squares solution.
- $b-Ax=b-\text{proj}_{\text{col}(A)}b$.
- $A^T(b-Ax)=A^T(b-\text{proj}_{\text{col}(A)}b)$.
- $\text{proj}_{\text{null}(A^T)}b=b-\text{proj}_{\text{col}(A)}b$.
- Thus, $A^T(b-Ax)=0$ or $A^T Ax=A^T b$.
- This is called the normal equation associated with $Ax=b$.

Theorem 7.8.3

(a) *The least squares solutions of a linear system $A\mathbf{x} = \mathbf{b}$ are the exact solutions of the normal equation*

$$A^T A \mathbf{x} = A^T \mathbf{b} \quad (11)$$

(b) *If A has full column rank, the normal equation has a unique solution, namely*

$$\hat{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b} \quad (12)$$

(c) *If A does not have full column rank, then the normal equation has infinitely many solutions, but there is a unique solution in the row space of A . Moreover, among all solutions of the normal equation, the solution in the row space of A has the smallest norm.*

- Proof: (a) done
- (b). Theorem 7.5.10 implies $A^T A$ is invertible.
- (c) omit.
- Example 3.

Error vector

- $\mathbf{b} = \text{proj}_{\text{col}(A)}\mathbf{b} + \text{proj}_{\text{null}(A)^{\top}}(\mathbf{b})$.
- $\mathbf{b} - A\mathbf{x} = (\text{proj}_{\text{col}(A)}\mathbf{b} - A\mathbf{x}) + \text{proj}_{\text{null}(A)^{\top}}\mathbf{b}$.
- By (7) $\text{proj}_{\text{col}(A)}\mathbf{b} - A\mathbf{x} = 0$ if \mathbf{x} is lss.
- $\mathbf{b} - A\mathbf{x}' = \text{proj}_{\text{null}(A)^{\top}}\mathbf{b}$.
- Least squares error = $\| \mathbf{b} - A\mathbf{x}' \|$
 $= \| \text{proj}_{\text{null}(A)^{\top}}\mathbf{b} \|$.

Theorem 7.8.4 *A vector $\hat{\mathbf{x}}$ is a least squares solution of $A\mathbf{x} = \mathbf{b}$ if and only if the error vector $\mathbf{b} - A\hat{\mathbf{x}}$ is orthogonal to the column space of A .*

- Example 4.

Fitting a curve to experimental data

- Data $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$.
- $y = a + bx$. Find the best a, b .
- If the line passes through the data, then
- $Mv = y$ where

$$M = \begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_n \end{bmatrix}, v = \begin{bmatrix} a \\ b \end{bmatrix}, y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}$$

- $M^T M v = M^T y$. Normal system
- If the x-coordinates are all distinct, then M has column rank 2 which is full,
- $v = (M^T M)^{-1} M^T y$.
- v gives us $y = ax + b$, the least squares line of best fit or regression line.
- What is minimized is $S = (y_1 - (a + bx_1))^2 + \dots + (y_n - (a + bx_n))^2$, the squares of residuals.
- Example 5.

Least squares by higher-degree polynomials

- Data $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$.
- $y = a_0 + a_1x + \dots + a_mx^m$. ($m < n-1$)
- Again, we can write this as:

$$Mv = y,$$

$$M = \begin{bmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^m \\ 1 & x_2 & x_2^2 & \cdots & x_2^m \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_n & x_n^2 & \cdots & x_n^m \end{bmatrix}$$

- If $m \geq n-1$, then exact solutions exist. (Lagrange interpolation formula)
- If $m < n-1$, we need to find the best solution.
- $v = (M^T M)^{-1} M^T y$.
- Example 7 to find the gravitational constant. (Read yourselves.)