


3.1. Operations on matrices


Matrix notation, operations, row and column vectors, product AB(important), transpose



Matrix notation

- ▶ Matrix: a rectangular array of real numbers.
- ▶ $m \times n$ -matrix: m rows and n columns
- ▶ square matrix: $n \times n$ -matrix
- ▶ Notation $A = [a_{ij}]_{m \times n}$, $A = [a_{ij}]$, $(A)_{ij} = a_{ij}$

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$



Operations on matrices

- ▶ $A=B$ if and only if they have same size and the same entries: $a_{ij}=b_{ij}$ for all i,j ,
- ▶ $A+B$: $(A+B)_{ij}=A_{ij}+B_{ij}$
- ▶ $A-B$: $(A-B)_{ij}=A_{ij}-B_{ij}$
- ▶ cA : $(cA)_{ij}=c(A)_{ij}=ca_{ij}$, $-A= (-1)A$.
- ▶ See Ex 1,2,3.



Row and column vectors

- ▶ Row vectors: $r = [r_1, r_2, \dots, r_n]$; i.e., $1 \times n$ matrix
- ▶ Column vectors: $c = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{bmatrix}$
- ▶ Think of $m \times n$ matrix as m row n -vectors in a column.
- ▶ Think of it as n column m -vectors in a row.




- ▶ $[c_1, c_2, \dots, c_n] =$

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} = \begin{bmatrix} r_1 \\ r_2 \\ \vdots \\ r_m \end{bmatrix}$$

- ▶ $r_i(A) = [a_{i1}, a_{i2}, \dots, a_{in}]$

- ▶ $c_j(A) = \begin{bmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{mj} \end{bmatrix}$



Define Ax where A: mxn-matrix, x n-vector

$$\begin{array}{l} \text{▶ } a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2 \\ \vdots \qquad \vdots \qquad \ddots \qquad \vdots \qquad \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m \end{array}$$

$$\begin{bmatrix} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

$$x_1 \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} + x_2 \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix} + \dots + x_n \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$




- We define Ax to be:

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = x_1 \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} + x_2 \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix} + \dots + x_n \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix}$$

- $Ax = [a_1, a_2, \dots, a_n] = x_1 a_1 + \dots + x_n a_n$
where a_i are column vectors.

Thus, we can write a system of linear equations
as $Ax = b$ for b $m \times 1$ column vector.



- ▶ See examples top. Page 83 and Example 4.
- ▶ Linearity Property: Theorem 3.1.5:
A $m \times n$ matrix, u, v column n -vector. Then
 - (a) $A(cu) = c(Au)$, c a real number
 - (b) $A(u+v) = Au + Av$.
 - Equivalently $A(cu+dv) = c(Au) + d(Av)$, c, d reals
- ▶ Proof: Simply use definitions and follow...



Product AB: Natural Definition

- ▶ A $m \times n$ -matrix, B $n \times r$ matrix. Define AB $m \times r$ -matrix so that $(AB)x = A(Bx)$ for any r -vector x .
- ▶ This is the associativity which is needed....
 - $B = [b_1, b_2, \dots, b_r]$
 - $Bx = x_1 b_1 + x_2 b_2 + \dots + x_r b_r$
 - $A(Bx) = A(x_1 b_1 + x_2 b_2 + \dots + x_r b_r)$
 $= x_1 A b_1 + x_2 A b_2 + \dots + x_r A b_r$
 - $(AB)x$ must equal $x_1(Ab_1) + x_2(Ab_2) + \dots + x_r(Ab_r)$
- ▶ Definition $AB = [Ab_1, Ab_2, \dots, Ab_r]$.
- ▶ See Example 5.



Specific entry of AB

- ▶ A mxs matrix, B sxn matrix -> AB mxn matrix
- ▶ $(AB)_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{is}b_{sj}$
 $= r_i(A)c_j(B)$ or $r_i(A).c_j(B)$ (second as vectors)
- ▶ Proof: $AB = [Ab_1, Ab_2, \dots, Ab_n]$
=



► Remove brackets to get a familiar formula

$$\left[\begin{array}{cccc} a_{11}b_{11} + & a_{12}b_{21} + & \dots & + a_{1s}b_{s1} \\ a_{21}b_{11} + & a_{22}b_{21} + & \dots & + a_{2s}b_{s1} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1}b_{11} + & a_{m2}b_{21} + & \dots & a_{mn}b_{s1} \end{array} \right], \left[\begin{array}{cccc} a_{11}b_{12} + & a_{12}b_{22} + & \dots & + a_{1s}b_{s2} \\ a_{21}b_{12} + & a_{22}b_{22} + & \dots & + a_{2s}b_{s2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1}b_{12} + & a_{m2}b_{22} + & \dots & a_{mn}b_{s2} \end{array} \right], \dots, \left[\begin{array}{cccc} a_{11}b_{1n} + & a_{12}b_{2n} + & \dots & + a_{1s}b_{sn} \\ a_{21}b_{1n} + & a_{22}b_{2n} + & \dots & + a_{2s}b_{sn} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1}b_{1n} + & a_{m2}b_{2n} + & \dots & a_{mn}b_{sn} \end{array} \right]$$

- Theorem, The ij entry of AB is the product of i -th row vector of A times the j -th column vector of B .
- See Example 7.



Finding rows and columns of AB.

- ▶ $AB = [Ab_1, \dots, Ab_n]$. Thus $c_j(AB) = Ac_j(B)$.
This is the column rule
 - $Ax = x_1c_1(A) + \dots + x_sc_s(A)$ (A $m \times s$ \times $s \times 1$)
- ▶ $r_i(AB) = r_i(A)B$. Row rule. (Use Dot Product rule to see this.)

$$AB = \begin{bmatrix} r_1(A)B \\ r_2(A)B \\ \vdots \\ r_m(A)B \end{bmatrix}$$

- $yB = y_1r_1(B) + \dots + y_sr_s(B)$ (B $s \times n$, y $1 \times s$) (To see, this, $yB = [yb_1, \dots, yb_n] =$

$$\left[\begin{bmatrix} y_1 b_{11} + \\ y_2 b_{21} + \\ \vdots \\ y_s b_{s1} \end{bmatrix}, \begin{bmatrix} y_1 b_{12} + \\ y_2 b_{22} + \\ \vdots \\ y_s b_{s2} \end{bmatrix}, \dots, \begin{bmatrix} y_1 b_{1n} + \\ y_2 b_{2n} + \\ \vdots \\ y_s b_{sn} \end{bmatrix} \right] = \begin{bmatrix} y_1 b_{11}, y_1 b_{12}, \dots, y_1 b_{1n} \\ + y_2 b_{21}, + y_2 b_{22}, \dots, + y_2 b_{2n} \\ \vdots \\ + y_s b_{s1}, + y_s b_{s2}, \dots, + y_s b_{sn} \end{bmatrix} = \begin{bmatrix} y_1 r_1(B) \\ + y_2 r_2(B) \\ \vdots \\ + y_s r_s(B) \end{bmatrix}$$

► Theorem 3.1.8.

- (a) the j-th column of AB is a linear combination of columns of A with coefficients from j-th column of B.
- (b) the i-th row of AB is a linear combinations of rows of B with coefficients from the i-th row of A.



Transpose

- ▶ A $m \times n \dots A^T m \times n$ rows become columns and vice versa.
- ▶ $(A^T)_{ij} = (A)_{ji}$.
- ▶ See Example 10.



Trace

- Given $n \times n$ -matrix A (1×1 also), $\text{tr}(A) = \text{sum of the diagonal entries } A_{11}, A_{22}, \dots, A_{nn}$.


$$A = \begin{bmatrix} 1 & 0 & 3 \\ -1 & -1 & 4 \\ 1 & 1 & 1 \end{bmatrix}, \text{tr}A = 1$$

- $\text{tr}A = \text{tr}A^T$.



Inner and outer matrix product

- ▶ Same size ($n \times 1$) column matrix u, v
- ▶ $u^T v$ is 1×1 matrix or a number: matrix inner product.
 - $u^T v = u \cdot v = v \cdot u = v^T u$
- ▶ uv^T is $n \times n$ matrix: matrix outer product
- ▶ See Example 11
- ▶ $\text{tr}(uv^T) = \text{tr}(vu^T) = u \cdot v.$




Ex set 3.1.

- ▶ 1–14 recognition, computations.
- ▶ 16–22 computations, rules,,, traces



3.2. Algebraic properties

Properties, inverses



Addition, scalar multiplication rules

- ▶ Theorem 3.2.1. a, b scalars, A, B, C same size
 - (a) $A+B=B+A$ commutativity
 - (b) $A+(B+C)=(A+B)+C$
 - (c) $(ab)A=a(bA)$
 - (d) $(a+b)A=aA+bA$
 - (f) $a(A+B)=aA+aB$
- ▶ Proof: simple calculations...



Multiplication rules

- ▶ AB is not necessarily equal BA . (not commutative) See Example 1.
- ▶ Some times $AB=BA$. Then A and B commute.
 - See Example *
- ▶ Theorem 3.2.2. $a, A \text{ mxn } B \text{ pxq } C \text{ rxs}$
 - (a) $A(BC)=(AB)C$ ($n=p, q=r$)
 - (b) $A(B+C)=AB+AC$. ($n=p=r, q=s$)
 - (c) $(B+C)A=BA+CA$. ($q=s=m, p=r$)
 - (d) $A(B-C)=AB-AC$ (e) $(B-C)A=BA-CA$
 - (f) $a(BC)=(aB)C=B(aC)$ ($q=r$)




- ▶ Proof (a) The rest is omitted.
 - Let c_j be the j -th column of C .
 - Question: What is j -th column of DC for some D ?
 - j -th column of $(AB)C$ is $(AB)c_j$.
 - j -th column of $A(BC)$ is $A(BC)_j$. $(BC)_j = Bc_j$. Thus, $A(Bc_j)$.
 - We showed $A(Bx) = (AB)x$ for any vector x .
- ▶ Zero matrix O : all the entries are 0.
- ▶ $A+O=O+A=A$; must be of same size
- ▶ $A-A=A+(-A)=O$
- ▶ $0A=O$
- ▶ $OA=O$ (may be different size zero matrices)
- ▶ If $cA=O$, then either $c=0$ or $A=O$.
- ▶ See Example 3. $AB=O$ but BA is not O .



Identity matrix.

- ▶ [1],
$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$
- ▶ I_m denotes $m \times m$ identity matrix
- ▶ A $m \times n$ matrix $A I_n = A$, $I_m A = A$
- ▶ Note that I_m is already in reduced row echelon form.
- ▶ Conversely, a row reduced $m \times m$ -matrix either has zero rows or equals I_m (Theorem 3.2.4)



Inverses

- ▶ A $n \times n$ matrix. If B is $n \times n$ and satisfy $AB = BA = I_n$, then A is invertible and B is an inverse of A .
- ▶ See Example 4.
- ▶ A matrix may not have an inverse. (See Example 5)
 - When a row or a column of it is zero.
 - When two rows (or two columns) are the same...
 - But there are more than these...
 - We will figure out precisely when...



Properties of Inverse

- ▶ Theorem 3.2.6. A invertible. B,C inverses.
Then B=C.
- ▶ Thus, we denote the inverse of A as A^{-1} .
- ▶ The inverse of 2x2-matrix is easy to obtain:


$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, ad - bc \neq 0, A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

- ▶ This is a computational result
- ▶ Example **, Example 8.




- ▶ $(AB)^{-1} = B^{-1}A^{-1}$, A,B $n \times n$ matrix
- ▶ Proof: $(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AA^{-1} = I$ and $(B^{-1}A^{-1})AB = B^{-1}(A^{-1}A)B = B^{-1}B = I$. Now use uniqueness

A product of any number of invertible matrices is invertible, and the inverse of the product is the product of the inverses in the reverse order.




Powers of matrix

- $A^0 = I$, $A^n = AA\dots A$ n-times
 - $A^{-n} = (A^{-1})^n = A^{-1}A^{-1}\dots A^{-n}$ n-times
 - $A^{r+s} = A^r A^s$, $(A^r)^s = A^r A^r \dots A^r = A^{rs}$
- **Theorem 3.2.9.** A invertible
- (a) A^{-1} is invertible and $(A^{-1})^{-1} = A$.
 - (b) A^n is invertible and $(A^n)^{-1} = A^{-n} = (A^{-1})^n$.
 - (c) kA is invertible and $(kA)^{-1} = k^{-1}A^{-1}$




- ▶ Proof (b): $A^{-n}A^n = (A^{-1})^n A^n$. $A^n A^{-n} = A^n (A^{-1})^n = I$ by cancelation from the middle.
- ▶ Thus, this operation is very similar to taking powers in real numbers....



Matrix polynomials.


- ▶ $p(x) = a_0 + a_1x + \dots + a_mx^m$
- ▶ Matrix polynomial in A ($n \times n$ -matrix):
 - $p(A) = a_0I + a_1A + \dots + a_mA^m$.
- ▶ See Example 12.
- ▶ $p_1(A)p_2(A) = (p_1 p_2)(A) = (p_2 p_1)(A) = p_2(A)p_1(A)$



Transpose again

Theorem 3.2.10 *If the sizes of the matrices are such that the stated operations can be performed, then:*

- (a) $(A^T)^T = A$
- (b) $(A + B)^T = A^T + B^T$
- (c) $(A - B)^T = A^T - B^T$
- (d) $(kA)^T = kA^T$
- (e) $(AB)^T = B^T A^T$



- ▶ Proof (e): A $m \times n$, B $n \times s$. AB $m \times s$ $(AB)^T$ $s \times m$
 - $B^T A^T$ $s \times m$ also.
 - ji -th entry of $(AB)^T$ is $(AB)_{ij} = r_i(A)c_j(B)$
 - $= r_j(B^T)c_i(A^T)$ is the ji -th entry of $B^T A^T$.
 - We need to show
 $r_i(A)c_j(B) = r_j(B^T)c_i(A^T)$

The transpose of a product of any number of matrices is the product of the transposes in the reverse order.




Properties of traces

Theorem 3.2.11 *If A is an invertible matrix, then A^T is also invertible and*

$$(A^T)^{-1} = (A^{-1})^T$$


Theorem 3.2.12 *If A and B are square matrices with the same size, then:*

- (a) $\text{tr}(A^T) = \text{tr}(A)$
- (b) $\text{tr}(cA) = c \text{ tr}(A)$
- (c) $\text{tr}(A + B) = \text{tr}(A) + \text{tr}(B)$
- (d) $\text{tr}(A - B) = \text{tr}(A) - \text{tr}(B)$
- (e) $\text{tr}(AB) = \text{tr}(BA)$



- ▶ 3.2.12 (e) state $\text{tr}(AB)=\text{tr}(BA)$ for square matrices. See Example.
- ▶ Proof: A $n \times m$, B $m \times n$ AB $n \times n \dots$

$$\begin{aligned}
 \text{tr}(AB) &= \sum_{i=1}^n (AB)_{ii} = \sum_{i=1}^n \sum_{j=1}^m A_{ij} B_{ji} = \sum_{j=1}^m \sum_{i=1}^n B_{ji} A_{ij} \\
 &= \sum_{j=1}^m (BA)_{jj} = \text{tr}(BA)
 \end{aligned}$$



- ▶ Product of row and column vector to be useful later

Theorem 3.2.13 *If \mathbf{r} is a $1 \times n$ row vector and \mathbf{c} is an $n \times 1$ column vector, then*

$$\mathbf{rc} = \text{tr}(\mathbf{cr}) \quad (11)$$

- ▶ Proof: Let $\mathbf{u}=\mathbf{r}^T$, $\mathbf{v}=\mathbf{c}$. $\mathbf{u}^T\mathbf{v}=\text{tr}(\mathbf{uv}^T)$. ((25) sec3.1)
 - Thus $\mathbf{rc}=\text{tr}(\mathbf{r}^T\mathbf{c}^T)=\text{tr}((\mathbf{cr})^T)=\text{tr}(\mathbf{cr})$



Transpose and dot product.

- ▶ $\mathbf{Au} \cdot \mathbf{v} = \mathbf{u} \cdot \mathbf{A}^T \mathbf{v}$ and $\mathbf{u} \cdot \mathbf{Av} = \mathbf{A}^T \mathbf{u} \cdot \mathbf{v}$
- ▶ Proof: Use $\mathbf{u} \cdot \mathbf{v} = \mathbf{v}^T \mathbf{u}$. -(*) Why true?
 - $\mathbf{Au} \cdot \mathbf{v} = \mathbf{v}^T (\mathbf{Au}) = (\mathbf{v}^T \mathbf{A}) \mathbf{u} = (\mathbf{Av})^T \mathbf{u} = \mathbf{u} \cdot \mathbf{Av}$.
 - $\mathbf{u} \cdot \mathbf{Av} = (\mathbf{Av})^T \mathbf{u} = (\mathbf{v}^T \mathbf{A}^T) \mathbf{u} = \mathbf{v}^T (\mathbf{A}^T \mathbf{u}) = \mathbf{A}^T \mathbf{u} \cdot \mathbf{v}$
- ▶ In the dot product, A moves across the dot by transposing....



Ex. Set 3.2.

- ▶ 1–6 confirmation by direct computations for specific matrices
- ▶ 7, 8 find unknown
- ▶ 9–12 confirmation
- ▶ 13–18 computations
- ▶ 32–37 a bit harder

