

7.2. Properties of Basis

Unique linear combinations

- If v_1, v_2, v_3 is dependent, then one can write a given vector v in more than one way as a linear combinations.
- Example $(1,1), (2,1), (3,2)$. $(6,4) = 2(3,2) = (1,1) + (2,1) + (3,2)$
- For independent set, this does not happen: If there are two linear combinations, then we subtract to get 0 vector written as a linear combination.
- This gives us “coordinates”.

Theorem 7.2.1 *If $S = \{v_1, v_2, \dots, v_k\}$ is a basis for subspace V of R^n , then every vector v in V can be expressed in exactly one way as a linear combination of the vectors in S .*

Removing and adding to get basis.

Theorem 7.2.2 *Let S be a finite set of vectors in a nonzero subspace V of \mathbb{R}^n .*

- (a) If S spans V , but is not a basis for V , then a basis for V can be obtained by removing appropriate vectors from S .*
- (b) If S is a linearly independent set, but is not a basis for V , then a basis for V can be obtained by adding appropriate vectors from V to S .*

- Proof: (a) Choose the second vector. If it is a linear combination of the previous vectors, then remove it. Otherwise, leave it alone. The result still span V . Continue to the next one until we cannot remove any. The result is independent.
- (b) Omit.

Theorem 7.2.3 *If V is a nonzero subspace of R^n , then $\dim(V)$ is the maximum number of linearly independent vectors in V .*

- Proof: If $\{v_1, \dots, v_s\}$ is independent with s maximum, then one cannot add any more vectors in V not in the span. This means V is the span of $\{v_1, \dots, v_s\}$.

Subspaces of subspaces: dimensions.

- V subspace of W , a subspace of \mathbb{R}^n .
- What can we say about dimensions?
- This follows since if there is a basis in V , we can add vectors in W , to obtain a basis of W . $\dim V \leq \dim W$
- If the dimension is the same, then the basis of V is the basis of W and $V=W$.

Theorem 7.2.4 *If V and W are subspaces of \mathbb{R}^n , and if V is a subspace of W , then:*

(a) $0 \leq \dim(V) \leq \dim(W) \leq n$

(b) $V = W$ if and only if $\dim(V) = \dim(W)$

- Some consequences

Theorem 7.2.5 *Let S be a nonempty set of vectors in R^n , and let S' be a set that results by adding additional vectors in R^n to S .*

- (a) *If the additional vectors are in $\text{span}(S)$, then $\text{span}(S') = \text{span}(S)$.*
- (b) *If $\text{span}(S') = \text{span}(S)$, then the additional vectors are in $\text{span}(S)$.*
- (c) *If $\text{span}(S')$ and $\text{span}(S)$ have the same dimension, then the additional vectors are in $\text{span}(S)$ and $\text{span}(S') = \text{span}(S)$.*

Theorem 7.2.6

- (a) *A set of k linearly independent vectors in a nonzero k -dimensional subspace of R^n is a basis for that subspace.*
- (b) *A set of k vectors that span a nonzero k -dimensional subspace of R^n is a basis for that subspace.*
- (c) *A set of fewer than k vectors in a nonzero k -dimensional subspace of R^n cannot span that subspace.*
- (d) *A set with more than k vectors in a nonzero k -dimensional subspace of R^n is linearly dependent.*

Example

- $v=(1,1,1), w=(2,1,-1), u=(2,0,1)$
- Basis of \mathbb{R}^3 .
- Write $(5,2,1)$ as a linear combination of v, w, u .

Theorem 7.2.7 *If A is an $n \times n$ matrix, and if T_A is the linear operator on R^n with standard matrix A , then the following statements are equivalent.*

- (a) *The reduced row echelon form of A is I_n .*
- (b) *A is expressible as a product of elementary matrices.*
- (c) *A is invertible.*
- (d) *$A\mathbf{x} = \mathbf{0}$ has only the trivial solution.*
- (e) *$A\mathbf{x} = \mathbf{b}$ is consistent for every vector \mathbf{b} in R^n .*
- (f) *$A\mathbf{x} = \mathbf{b}$ has exactly one solution for every vector \mathbf{b} in R^n .*
- (g) *$\det(A) \neq 0$.*
- (h) *$\lambda = 0$ is not an eigenvalue of A .*
- (i) *T_A is one-to-one.*
- (j) *T_A is onto.*
- (k) *The column vectors of A are linearly independent.*
- (l) *The row vectors of A are linearly independent.*
- (m) *The column vectors of A span R^n .*
- (n) *The row vectors of A span R^n .*
- (o) *The column vectors of A form a basis for R^n .*
- (p) *The row vectors of A form a basis for R^n .*